

## **An Economic Analysis of Cattle Numbers and Feed Sources in the USDA Basin and Range Region**

Report prepared for the Public Lands Council, Washington, DC

Stanley R. Johnson, Michael D. Helmar, Thomas R. Harris and Michael H. Taylor

University Center for Economic Development (UCED) Department of Economics  
and  
College of Agriculture, Biotechnology, and Natural Resources

University of Nevada, Reno  
Reno, Nevada  
(775) 784-1907  
[www.unr.edu/business/research-and-outreach/uced](http://www.unr.edu/business/research-and-outreach/uced)



This publication, *An Economic Analysis of Cattle Numbers and Feed Sources in the USDA Basin and Range Region*, was published by the University of Nevada Center for Economic Development. Funds for the publication were provided by the Public Lands Council, Washington, DC, through the National Center for Food and Agricultural Policy, Washington, DC. This publication's statements, findings, conclusions, recommendations, and/or data represent solely the findings and views of the authors and do not necessarily represent the views of the University of Nevada, or any reference sources used or quoted by this study. Correspondence regarding this document should be sent to:

University Center for Economic Development  
University of Nevada, Reno  
Department of Economics  
Mail Stop 204  
Reno, Nevada 89557



UCED  
University of Nevada, Reno  
Nevada Cooperative Extension  
Department of Economics

# **An Economic Analysis of Cattle Numbers and Feed Sources in the USDA Basin and Range Region**

by Stanley R. Johnson, Michael D. Helmar, Thomas R. Harris and Michael H. Taylor\*

[drstanleyjohnson@gmail.com](mailto:drstanleyjohnson@gmail.com)

## **1.**

### **Introduction**

Ranchers, rural communities, and those interested in agricultural production and the incidence of wildfires in the Basin and Range Region at present have little sound analysis to support related issues with the Bureau of Land Management (BLM), or other federal and state departments that fashion regulations and policies. These citizens and groups lack significant input into such regulations and policies. The result is that the many constituents that have a stake in the policies and regulations of federal and state lands are left out of important decisions about grazing, wildfires, wildlife habit and other major decisions about management of resources that impact their livelihoods.

This analysis is designed to provide those who have not had access to systematic information about rangeland issues mentioned above with information that can give them a substantive role as stakeholders. Examples of issues which could use the input of citizens, communities and agencies that are now left out of policy and regulatory decisions include:

How to deal with horse populations,

Differences between grazing perennials and invasive grasses,

Wildfire management and control,

Grazing after wildfires,

Active ranching versus allocations to wilderness, and

Other emerging policy and regulatory matters

One of the cardinal features of public policy and regulatory decisions is full inclusion of stakeholders. And, in regions where the federal and state governments control large areas of public lands there is reason to provide stakeholders analysis so they have a better informed voice in decisions than are now made.

---

\* Stanley R. Johnson is Assistant to the Dean, Michael D. Helmar is Research Manager, College of Agriculture, Biotechnology, and Natural Resources, Thomas R. Harris is Professor and Director of the University Center for Economic Development, Department of Economics, and Michael H. Taylor is Assistant Professor/State Specialist, Department of Economics/Cooperative Extension Service, all at the University of Nevada, Reno, Reno, Nevada 89557.

This report provides the beginnings of a stronger basis for broad participation and input of citizens, groups and private and public agencies into the decisions. The preliminary analysis provided in this report and perhaps other systematic analyses provides an opportunity for stakeholders to get their feet under the table in public land decisions that impact them. We wish to stress that the analysis is preliminary, and much of it is dependent on previously available scientific reports. For example, the costs and returns budgets from the land grant universities (unfortunately sometimes dated), the wildfire model that is only for a part of the Basin and Range Region, community economic models are just for Nevada, cheatgrass coverage estimates that are not complete, BLM and Forest Service AUM numbers that are not completely aligned with NASS cattle numbers, and the scoring analysis that at this point may be incomplete. Pinion and juniper encroachment is not addressed. But with these and other qualifications we have a start on getting more rancher and other citizen inclusion into range land decisions of federal and state agencies.

The focus of this analysis is two-fold: why have the cattle numbers been reduced in recent years and engaging in the current debate about cheatgrass grazing especially during the fall term of the year when it is edible by cattle. Of course, there are other issues that deserve input by the broad set of stakeholders. But we will concentrate on just two in this analysis, although the data summarized and presented can be used for a number of related issues.

## **2. Objectives**

The objectives of this research project will support several versions of the to-date untested policy and regulatory decisions about cheatgrass and reduced cattle numbers. The general objective is to provide the ranchers, rural communities and others stakeholders with vested interest in capacities to supply input into the decisions of the BLM on policy and regulatory decisions guiding public grazing lands management in the Region. It is high time that all of the participants in decisions about the range and the landscape in the Basin and Range Region have a full opportunity to provide input into the decisions that are now being more or less handed down from the federal agencies and in fact, several state agencies as well. Again, the general focus of our analysis is cattle numbers and cheatgrass grazing.

Specific objectives of this research project are the following:

Generation of complete costs and returns information (budgets) for counties in the Basin and Range Region. The information about the costs and returns has not been well maintained by the land grant universities in the selected states. We needed to up-date past versions of these costs and returns to begin our analysis about the range and landscape in the Basin and Range Region using concrete information on costs and returns. We needed variable costs and returns estimates by county and state for the Region to provide estimates of rancher net returns from cattle grazing. These “short term” estimates of returns can be input in estimating total revenue from allocations public lands to grazing, providing information for questions and discussions to promote a better understanding of the policies and regulations of the BLM.

Cattle numbers are the next step in defining grazing allotments and related issues in public policy and regulatory decisions for the Basin and Range Region. These cattle number (for beef cows) have been integrated together with the costs and returns information for counties in the Basin and Range Region. With these cattle numbers we will have full sets of estimated returns by county, state and for the Basin and Range Region. We can trace over time how these allotments of cattle numbers have changed and raise questions with the BLM about why these changes have generally moved downward and questions about addressing cheatgrass. These cattle numbers are from the USDA's National Agricultural Statistics Service (NASS), a federal agency that is assigned responsibility for providing data on county, state, and national agricultural production, including for beef cow and other cattle numbers.

We will compare data for BLM and Forest Service animal unit months (AUMs) with the NASS cattle numbers. The cattle numbers are generally available on a county basis, allowing bottom up aggregation for the Basin and Range regions in each state. However, the BLM and Forest Service AUM data is by state. In states that are not completely contained in the Basin and Range region, it is therefore difficult to compare cattle numbers directly with AUMs. Furthermore, even in the Basin and Range region, there are some states that have a substantial number of cattle that are grazed on private lands.

Although the primary focus of this study is on BLM grazing, in some states Forest Service numbers of AUMs are significant, and in some cases are greater than BLM AUM numbers. Considering both when comparing to cattle numbers gives us a more accurate view of public land allocation to cow-calf industries in various states. These sets of AUM data are from the Annual Reports available from the BLM and Forrest Service. In addition, the Forest Service data includes numbers of animals grazed by state on Forest Service lands. These data bases for AUMs will support a beginning set of estimates designed to generate comparisons to the NASS beef cattle numbers, and where they are different from the BLM (and Forrest Service), and how they are related to historical decisions for cattle numbers allocated to range lands. We will provide three sets of appendix tables (subsequently defined) to begin to sort out these questions.

In this research report our estimation of economic impacts of grazing on communities and states were confined to the State of Nevada and its counties. We selected Elko County in Nevada, which has seen a reduction in beef cow inventories by about 1/3<sup>rd</sup> compared to 1982. We then compared 1982 to 2016 as a basis for understanding the reduction decisions for allocation numbers by BLM. We also make an estimate of the state of Nevada which has many other larger industries than cattle grazing. This is an attempt to show how counties with substantial cattle numbers are disadvantaged by state level input-output analysis. These cattle numbers are indicative of an over simplification that is implied in evaluating results of county based input-output analysis where cattle are dominant.


We also included wildfire estimates in this analysis. But the wildfire analysis is only for a part of the Basin and Range Region—Nevada, Utah, Idaho and Oregon. The wildfire model was “run:” for the historical period of record. But this historical record is not reflective of the extent to which cheatgrass and other invasive species have moved into the Basin and Range Region. More recent data would include more area in cheatgrass because of recent wildfires that have permitted cheatgrass to dominate the perennial vegetation. In short, cheatgrass is becoming more and more a factor in rangeland allocations and management by BLM. This is because the BLM presently does not count cheatgrass in the grazing allotments. We will investigate this problem since it has been proven that cheatgrass can be grazed in the fall season, and at little cost to the perennials.

Finally, we will take up scoring of BLM range lands as a basis for informing improved decisions about cattle grazing, cheatgrass, and other range conditions in the Basin and Range Region. The calculated scores will pit the welfare of states, communities and wild life against increased cattle grazing and adjustments for cheatgrass allocations. How do cattle grazing allocations influence the welfare of communities and the state? Here we will take into consideration community and state damages of several types, smoke and wildlife reductions which affect the state and rural communities. Also the community analysis will include wild life habit.

This was a major task, using the available budgets, cattle numbers, community and state economic impact analysis, and wildfire data for our research. But it is important to obtain the data that reflect the nature of the problems of BLM in regulations, policies and management of cattle grazing on public lands. The analysis can be improved with additional resources, but it is at this point comprehensive, taking into consideration all of the facets of the problem and making sure that ranchers, rural communities and others interested in rangeland regulation and policy can be at the table when decisions about public land allocations are made.

### **3. Design of the Analysis**

The schematic design for the first part of the research project for the cattle grazing, state and community economics and wildfire analysis is provided in Figure 1. In a second Figure we provide a schematic sketch of the scoring approach to be applied in this analysis. Reviewing Figure 1, notice that it is related to budgets, cattle numbers, communities and wildfires.




Figure

### 1. Schematic Representation of the Economic Analysis of the Research Proposal

Also note from Figure 1, we have provided for a separation between the costs and returns which both feed into the cattle or beef cow numbers and community analysis. These elements of Figure 1 are prepared at the budget study geographic level which can then be aggregated to state totals and to Basin and Range Region totals. These totals illustrated in Figure 1 are however only for Basin and Range counties. That is, counties were omitted if not included in the Basin and Range region. This will be shown later in the report.

As well, the wildfire impacts in Figure 1 are shown to affect the cost side of the calculations to a greater extent than the returns side. This is because the cost is where the wildfire impacts are felt and reported. The revenue side is from the sales to the markets in the states and Basin and Range Region and has an impact only when the cattle and calves are sold from the ranches. The costs and revenues are as well related to states and community impacts on residents. These distinctions will be important as we proceed through the substance of the research report.

Figure 2 contains the elements that we will consider for the scaling analysis, and is shown below.



**Figure 2. Possibilities for Scoring in the context of BLM Lands**

Figure 2 is to support an application of scoring to better understand the motives of BLM and bringing in cattle numbers, communities, wildlife, and cheatgrass grazing. The way Figure 2 is structured is to include consideration of adding to the cattle numbers by opening and authorizing cheatgrass grazing in the fall after frost, when cheatgrass, can be grazed. The grazing of cheatgrass can last from early October until late December. If it is 2 months there are distinct advantages of including in the BLM allotments cheatgrass grazing. First, the grazing of cheatgrass does not require hay which would be fed to cattle during the 2 months during the fall season. This is a benefit to ranchers since they do not have to supply feed for their cattle taken off of the grazing range. Second, the grazing of cheatgrass is beneficial since it cleans up the current year's growth and the thatch from last year or previous years that is available. This reduces the fuel load and thus wildfire risk and intensifying subsequent years. Third, these benefits go to states and rural communities subject to recurring wildfires. Of course there are other benefits but these will be the ones selected for the scoring experiment.

A simple scoring equation will suggest this scoring approach. That is,

- 1) Increased Cattle numbers (+) Degrading of range land (-) Fuel load reduction (+) Increased wildfire retardation (+) Community wildfire protection (+) and Improved wildlife habitat (+-).

The idea is to put numbers or weights on the terms in this equation and add them up for supporting decisions about the BLM grazing allocations. Of course, this scoring mechanism must be for a particular area of land—presumably including rural communities and available wildlife habitat. But the other features of the scoring analysis can be easily transported by taking into consideration the calculation of the cheatgrass available for increasing the duration of cattle grazing appropriately and proceeding for the areas selected. As is clear from the equation, there are a number of terms that should be taken into consideration make the equation work for most areas.

There are a number of added BLM conditions that go with this scoring equation. In the areas, there should be scouting for the cheatgrass spring and summer growth that would be available for grazing. Some summers will be low growth and some summers will be high growth. We suggest some type of compromise related to these considerations. We decrease the cattle grazing somewhat during higher growth years than appropriate (for good years) and manage appropriately low years of the cheatgrass growth. In short, it takes active management to handle cheatgrass grazing levels in BLM managed lands. Still another possibility for low growth years would be to take the cattle off the cheatgrass earlier making them demand more feed, but at the same time managing the cheatgrass available on the land.

With these considerations we can proceed to discuss the components of the project more definitively.

The strategy for this project is to proceed from costs and returns (budgets) on a per cow estimated basis for counties to total cattle numbers on the basis of NASS estimates for the counties that are within the Basin and Range Region. This will provide us with *variable* net returns to total cattle grazing for each county. From these estimates, we can move to apply input–output models to provide for each county estimates of gross economic impact reflecting the inter-industry flows within and out of the counties. We used this input-output analysis for Nevada and for only one county and the full state for this analysis. The objective was to show the impact for Elko County compared to the full state impact. What is shown is the major impact that is estimated for Elko County against the impact for the state. The bias of state input-output models applied to states is clearly observed.

The next effort is to apply the wildfire model for a subarea of the Basin and Range Region to estimate the incidence of wildfires within this Region. This involves some manipulation since wildfires are more predominant currently than in the past. Several approaches were applied to perform an update the wildfire model. Then the model is applied for counties within the region to add to the results from the input-output analysis. Thus, we have from the wildfire model impacts for counties in the area observed as well as economic impacts for the state and counties. For the areas observed, the growing importance of wildfires is a factor in the allotments for cattle grazing as well for wild life and other environmental factors. We will take make estimates and try to obtain a consensus on the impacts of wildfires.

Finally, we will apply scoring models to evaluate the general cattle grazing, wildfires, and wild life habit for improving the economic welfare states and communities or the welfare of all constituents in the region. Again several versions of the scoring model will be tried as a basis for generating consensus results for the communities and the states in the Basin and Range Region. This will provide those not in the loop about BLM decisions a basis for arguments about systemic change in BLM policies and regulations.

Costs and Returns or Budget estimates

The analysis of costs and returns is for the Basin and Range Region which stretches from the states of Washington to Arizona from North to South and from the states of California to Wyoming from West to East. The first task was to prepare on a per cow basis variable costs and returns (budgets) available for various counties within the Basin and Range Region. This is obviously a rather large region, but fortunately within it the cattle budgets are rather similar in terms of technology. At the same time the budgets are often prepared for different years and often dated. The source of these budgets is the faculty and staff of land grant universities, and in fact, they are produced in various years. (California and Colorado were eliminated from this analysis due to the fact that there were no budgets available from these two states.) Thus, for this analysis, these budgets were “updated” to reflect the current period (2016). Additionally, in order to make the budgets and eventual aggregations more meaningful under general conditions, an average of costs and returns from 2014-2016 was used. This methodology was approved by the Budget Committee which included land grant academics from Utah, Idaho, Oregon and Nevada (affiliations are noted in appendix A).

### Cattle Numbers

Additionally, the averages of the cattle numbers in the counties and/or the states for the 2016 and the three year averages were utilized in the aggregation phase of the analysis. Cattle numbers came from the USDA National Agricultural Statistics Service (NASS) and are available for each state and counties within the Basin and Range Region. Cattle numbers are also available from the Census of Agriculture, as well. For those counties which cattle numbers were not available due to proprietary conditions, the NASS survey data and the Census data were used to extrapolate yearly county cattle numbers. For example, the latest Census year was 2012 (available in 2014) which provided us with cattle numbers in the reference year for all counties. Cattle inventories for counties in which the cattle numbers were not available in the annual NASS survey data were all extrapolated for 2015 and 2016. The Agricultural Statistician in Nevada provided these cattle numbers for all states in the Basin and Range Region. We selected the USDA/NASS for cattle numbers due to the fact that the BLM and Forest Service Districts often did not have these numbers for years needed to support the analysis or had only AUMs for the years included. We will review the BLM state numbers in a subsequent section

### County and state level assessments of economic impact

The economic assessments of counties and the state were based on input output models and for Nevada only. This is because the input output models have not been developed for other counties or states in the Basin and Range Region. We simply used a selected county available analysis for this report, Elko. The county has about 1/3<sup>rd</sup> of the cattle members in Nevada. But, these Nevada county numbers for economic impacts will be sufficient to give an idea about the “benefits” of cattle grazing in the state and in an agricultural county. In short, we used input output models for the State of Nevada and for the selected county in Nevada for this portion of the analysis. For the state, the cattle numbers estimates are not significant in the analysis for the economic input-output. Other industries are more economically important. But for the agricultural county, Elko County, they were quite significant. In recent years and for historical cattle numbers which have been reduced significantly? We wanted to get an approximation of how the approximate 1/3 greater cattle numbers of the past would impacted the economies of this agricultural county. For this comparison we used the 1982 Census year estimates of cattle numbers for the analysis of differences in cattle numbers. We will use this county estimate to

provide rough estimates of the economic impacts of increasing cattle numbers for other counties in the Basin and Range Region.

## Wildfires

Wildfires are closely linked to the prevalence of cheatgrass in the Basin and Range Region. For the analysis of wildfires, we employed a wildfire model for a subset of counties in the Basin and Range Region. This model covers the northern part of Nevada, most of the Basin and Range Region in Utah and the western parts of Oregon and Idaho. It is a model in which we have confidence and will serve as an illustration for the rest of the Basin and Range Region. However, here there is a possible problem with the historical period of record for which the wildfire model was initially run. We will apply to the Basin and Range Region wildfire model which has sufficient “granulation” that we can obtain estimates of the cheatgrass (and other invasive species) covering the land in the study area. Specifically, by investigating the period of historical record we can estimate the incidence of impacts of cheatgrass for wildfires.

Alternative approaches for application of the wildfire model can be undertaken in the last several years. First, we can simply use the historical period of record for the analyses. But this would be assuming that cheatgrass and other invasives have not increased in coverage of the land during the historical period. A second and more appropriate method is to use select current years (about 10) of land cover or an average of current years, and then apply the weather data from the historical period. This will provide an improved of the estimates of the impact of cheatgrass cover of land for the counties analyzed. We may as well attempt obtain estimates from the BLM officials about the cheatgrass cover in the counties. If successful, we will use these estimates as well for comparison by running separate versions of the wildfire model. How we will estimate cheatgrass cover of land will be used to provide a probability of the incidence of wildfires. It is well known that cheatgrass wildfires occur every 5 – 8 years and that wildfires in the perennials occur every 50 – 80 years. Thus, with sufficient granulation we can estimate the incidence of fires from cheatgrass from the historical and more current years of data.

## Scoring related to wildfires and economic criterion

Scoring is related to determining the best alternative when there are several competing factors for the public land. In this case we will utilize scoring to reflect the competing interests related to grazing on public lands managed by BLM on one hand and the economic impacts for the state and rural communities, late season cheatgrass grazing possibilities, wild life habit for the rural communities. That is, late season grazing and increased cattle numbers will be featured in the scoring analysis. We know that cattle can graze cheatgrass during the fall season compared to the spring season when the grazing is unpredictable. This grazing period is not preferred by ranchers due to the shortness of the spring season and the unpredictability of when the spring grazing season will be available. Spring is a season with many factors that explain the grazing period.

We also know that cheatgrass can be controlled by late season grazing after frost when the burrs on the cheatgrass and other invasives are no longer harmful and are edible for cattle. This provides a natural opportunity for an assessment of cheatgrass control. That is, during the fall cheatgrass can be controlled by the cattle on it from about October 1 removing cattle when it begins to snow, in mid-December. With small amounts of protein supplement made available, the cattle will consume the cheatgrass from the current year and as well consume “thatch” from previous years, with minimal

grazing of perennials. We have experience to show that the cheatgrass grazing does not diminish the gains or condition of cattle.

Protein supplement can be dispensed from a feeder with the feeder pulled along in the area where the cattle are grazing occurs-- about 200-300 yards from the feeder. The scoring will be applied to compare areas with cattle grazing which leads to a reduction of wildfires that occur with a reduced build up for fuel. This will involve a wildfire model with probabilities of wildfire managed with and without cheatgrass grazing. It may be for instance, that BLM could afford to provide credits to cattle grazing on cheatgrass in the fall. These credits could be for a reduction in the stipend that is generated from cattle grazing during the permitted season---a one to one trade off. The result will feature reduced cheatgrass from annual and hold over thatch, and benefits to states and communities through increased cattle grazing, increased cattle numbers, economic impact analysis and wildlife habitat along with results of potentially decreases in the range environment. Several versions of the scoring rules will be developed and presented for evaluation.

#### **4. Budgets for Cattle Grazing, in the Basin and Range Region**

Two sets of budgets were prepared that include all counties in the Basin and Range Region (with California and Colorado omitted). These budgets were for 2016; and average of the years 2014, 2015 and 2016. The corresponding cattle numbers were averaged for the three-year period as well reflecting the cattle numbers in the 2014 - 2016 years. In the Basin and Range Region there were 169 counties required for inclusion in the budgets or costs and returns. Some of the budgets are for specific counties and some collections of counties. Again, we used the budgets prepared by land grant institutions in each state to define the specific counties or collections of counties for application of the estimates. We used the three year averages as per agreement from our Budget Committee that 2016 was an off year for cattle production and sales due to both low prices and low cattle numbers

The counties within the Basin and Range Region for each state are reported in Appendix B (in graphic and listed form). Note that there are tables and/or figures for each state; Washington, Idaho, Montana, Wyoming, Utah, Oregon, Nevada, New Mexico and Arizona. California and Colorado have some counties in the Basin and Range Region but did not have available budgets for our study. Colorado had only a stocker budget and California did not have available budgets for counties or portions of counties in the Basin and Range Region. Thus in this analysis we omitted these states and counties from the calculations of variable costs and returns. The counties in the omitted states could be included, if budgets were available or we could use the budgets from bordering states for application.

A major problem in developing the budgets was the fact that the land grants in the Region had not kept up with the changing times. Because most of the budgets were not for 2014, 2015, and 2016, they needed to be updated to reflect the years for analysis. The process that we used for updating involved several sets of available numbers, including State Agricultural Statistics, the USDA National Agricultural Statistics and Prices Paid by Ranchers state data, and Bureau of Labor price indices. We in fact, used a combination of these data series to update the budgets. For the prices of cattle, we used the USDA reported Commodity returns from cow-calf Operations in the Basin and Range Region, and for hay and other cost requirements we used the Prices Paid from the states and other price indices that were available from the Bureau of Labor Statistics. This approach was endorsed by the Budget Committee.

The budgets for 2014–2016 are in Appendix C and the budgets for 2016 alone are in Appendix D. Note that the numbers at the top of the budget pages provides information on the year (or years) when the budgets were actually prepared by the state land grant institutions for the respective states. Again, some these are for at least 10 years older than for the ears of the analysis. Also note again that the updated budgets were prepared only for variable costs and returns. We decided to leave the long term “fixed” costs out of the budget analysis because they do not reflect the short-term annual decisions made by ranchers for production. Variable costs and returns are in fact, the appropriate numbers for this analysis.

Reviewing the budgets across states, we found that the basic technologies were more or less the same for cow-calf or cattle production. The primary differences in bred cow costs and returns were due to efficiencies gained from scale of operation, growth conditions for forage and climate, and the proportion of feed supplied when grazing occurred on public lands. Revenues per bred cows and calves for 2014-2016 range between \$540 and \$1,253 across the region and variable costs ranged from \$147 to \$1,106. The variable cost estimates that stick out are for the counties and collections of counties in the states of Arizona on the low side and Oregon on the high side. The lowest overall revenues were for Arizona, with the highest in the Washington area, but Idaho and Montana also yielded higher than average revenues per cow.

## **5. NASS Cattle Numbers**

The USDA National Agricultural Statistics Service (NASS) “beef cow” inventories are available for each county within the Basin and Range Region. These numbers or “estimates” however require some qualification. The actual numbers are available for each county during the Census years, and had to be extrapolated for non-Census years, except for counties with large beef cow numbers where they were directly available from NASS (but also on an interpolated basis). With this qualification, the numbers were used and available for all of the BLM counties. For the three year budgets presented as summarized in the previous section, the beef cow numbers had to be averaged as well. These county averages are presented in Appendix E. Again, we felt that the numbers or estimates for a three-year period would be better for use than 2016 numbers alone in the development of beef cow costs and returns for the Basin and Range Region.

The beef cow numbers however require added explanations. Beef cows are provided for the year and lead to cow calf production. These numbers are factored or adjusted by NASS to account for replacements and or culling of beef cows. See the NASS explanation for the reference (need reference). The beef cows are put on the grazing range land mostly after calving and the calves are sold at 400+ pounds live weight. This is the major source of revenue for the beef cow growers, even when culling of cows is included. These beef cow numbers are the ones for use for evaluating costs and returns for the cattle operations in counties.

With the three- year average beef cow inventory numbers and the three year averages for the budgets, we can compute total variable costs and returns for each county and state in the Basin and Range Region. The results for the budgeted counties or regions are provided in Appendix F. The state estimates for sets of counties (or single counties) in the Basin and Range Region are summarized in Table 1.

First the total net variable returns (variable returns less variable costs) for the Basin and Range Region for the average period 2014-2016 were 1,202,220,000 or about 1.20 billion dollars. This total was distributed by state for BLM lands (which are often only a part of the stats) as shown in Table 1. Montana had the highest beef cow inventories (767,245) followed by Utah and Oregon at approximately 329,117 and 329,667, respectively. In county or collections of county numbers the largest beef cow numbers were in Elk County, Nevada, Klamath/lake Counties Oregon and Southwest New Mexico, all in the neighbor of 72,000 plus. Smaller numbers of beef cows on range land were mostly in the Nevada counties. More refined numbers are available in the county estimates in Appendix Table F and on the regional levels on Table 1.

**Table 1. Total Revenues, Variable Costs, and Returns**

	Thousand dollars			
	Beef Cow Inventory	Total Revenue	Total Costs	Returns
<b>Arizona</b>				
Plateau Region	35,106	\$21,987	\$9,828	\$12,158
SW Desert Region	40,710	\$21,998	\$5,983	\$16,014
Strip Region	13,498	\$8,302	\$6,654	\$1,648
Western Desert Region	50,542	\$27,436	\$10,671	\$16,765
Arizona Basin & Range	139,856	\$79,722	\$33,136	\$46,586
<b>Idaho</b>				
Northern Idaho	42,294	\$49,414	\$29,848	\$19,566
Magic Valley	94,333	\$97,512	\$47,749	\$49,762
Eastern Idaho	142,784	\$166,774	\$89,139	\$77,634
Idaho Basin & Range	279,411	\$313,699	\$166,737	\$146,963
<b>Montana</b>				
Montana Basin & Range	767,245	\$868,834	\$362,476	\$506,359
<b>New Mexico</b>				
Northwest New Mexico	37,833	\$30,396	\$12,930	\$17,466
Central New Mexico	44,633	\$35,064	\$13,852	\$21,212
Southwest New Mexico	79,491	\$59,336	\$23,268	\$36,068
New Mexico Basin & Range	161,957	\$124,797	\$50,061	\$74,736
<b>Nevada</b>				
Douglas County	5,653	\$5,168	\$3,376	\$1,792
Elko County	72,214	\$67,710	\$37,187	\$30,523
Eureka County	11,907	\$11,945	\$6,892	\$5,053
Humboldt County	36,597	\$35,901	\$22,763	\$13,137
Lyon County	11,453	\$10,695	\$7,224	\$3,471
Pershing County	11,887	\$11,621	\$6,841	\$4,780
White Pine County	15,561	\$17,382	\$10,768	\$6,614
Other Nevada Basin & Rang	57,763	\$56,068	\$33,221	\$22,847
Nevada Basin & Range	223,035	\$216,488	\$128,272	\$88,216
<b>Oregon</b>				
Klamath/Lake Counties	73,833	\$79,915	\$69,810	\$10,105
North Central Plateau	48,652	\$44,319	\$30,889	\$13,431
High Desert Area	197,532	\$174,004	\$156,164	\$17,840
Oregon Basin & Range	320,017	\$298,238	\$256,862	\$41,376
<b>Utah</b>				
Box Elder County	33,500	\$32,409	\$18,950	\$13,458
Duchesne County	25,000	\$26,371	\$10,966	\$15,405
Tooele County	13,767	\$13,711	\$4,996	\$8,715
Rich County	29,500	\$28,622	\$11,781	\$16,841
Utah Other Basin & Range C	227,900	\$226,805	\$104,566	\$122,239
Utah Basin & Range	329,667	\$327,917	\$151,259	\$176,658
<b>Washington</b>				
Washington Basin & Range	49,579	\$62,113	\$24,611	\$37,502
<b>Wyoming</b>				
Wyoming Basin & Range	49,579	\$299,424	\$215,599	\$83,824
<b>Total Basin &amp; Range</b>	<b>2,320,346</b>	<b>\$2,591,233</b>	<b>\$1,389,012</b>	<b>\$1,202,220</b>

Clearly, the numbers of beef cows available for grazing on the BLM lands could be increased, if wildfire and cheatgrass and other invasives were included in the basic calculations for authorized grazing numbers approved for the respective counties. Cheatgrass grazing is now not permitted in calculations of BLM lands for the Basin and Range Region. If authorized by the BLM, the numbers of beef cows grazed could be increased by roughly the amount of available land estimated as cheatgrass cover. The increase in beef cow numbers and the grazing of cheatgrass, in association with appropriate post-grazing vegetation restoration prescriptions could lead to reductions in wildfire incidence in the Basin and Range Region. How much of the cheatgrass that is potentially available will be estimated in the application of the wildfire model in the section below. This is a real question for the “BLM and management”—how much could the wildfire incidence be reduced, if additional beef cow numbers were authorized and cheatgrass grazing allowed. Again, this calculation has important implications for the Basin and Range Region as indicated in Equations 1. Also we mention that the improvement in habitat is important for sage grouse restoration, for example.

The next section is devoted to first tracking the BLM cattle numbers or AUMs, Forest Service numbers of AUMs and a comparison or consolidation to our beef cow numbers.

## **6. BLM Cow Calf AUMs and Beef Cow Comparisons**

The AUM numbers in this analysis are from three sources: the BLM data is from the Rangeland Administration System (RAS) for all states in the Basin and Range Region (Appendix G), the Forest Service AUM and grazed cattle data is from the U.S. Forest Service, *Grazing Statistical Summary* for all states in the Region (Appendix H), and the total Federal AUMs are an aggregation of the data from these two reports (Appendix I). In this section we will take each of these appendices in order and attempt to explain their implications for this study. As a first observation, we note that the explanations in Appendix G and H are in animal unit months (AUMs) and not identical in definition to the beef cow numbers from NASS. Thus the first approach is to attempt to reconcile these AUMs with beef cow numbers. It is very confusing to have several agencies reporting numbers that cannot be completely reconciled with other data sources. This issue needs careful attention when using and interpreting this information. With this qualification we will begin to provide an explanation of the AUMs reported by the BLM and Forest Service.

### BLM Reports on AUMs in the Selected States (Appendix G)

In Appendix G we find several measures of AUMs and beef cattle numbers. For the BLM numbers we have plotted in the charts “Active”, “Authorized” and “Active Less Suspended” AUMs for the historical record for the included states. Active AUMs are those that could be authorized for use on public lands. Authorized AUMs are our best information about the available AUMs for grazing consumption as they are those AUMs that are actually allowed for public use. There is some controversy about the suspended category since it includes several reasons for reducing the active category, and does not actually reflect the AUMs that are not in use. In other words, it falls far short of measuring the difference between active and authorized AUMs. We will use the authorized category for this analysis but there is

much to learn about the active less suspended category. This category needs better explanation from the BLM than is now available.

If we take the BLM authorized AUMs and compare them to beef cow numbers available for each state, we get varying results. For example, for Nevada where the Forest Service and other alternative sources of grazing land have relatively little land allocated to grazing, the ratio of cattle numbers from NASS equates to about five animal units per cow on average, or about five months of grazing per cow for the state. This compares to a roughly six-month grazing season for many of the operations in the state. That is if we use about six AUMs for cattle we get beef cow numbers that are roughly consistent. On the other hand, for Arizona and Utah, where the Forest Service has a substantial presence on potential grazing lands, we get very different numbers for beef cattle numbers compared to the BLM AUMs. The reason for this is that the Forest Service controls significant tracts of land in Arizona and Utah. Thus we need to capture the Forest Service AUMs before making such comparisons.

This set of comparisons is suggestive of still another possibility for land available for grazing, private lands. Many private lands are used for haying to prepare for the winter season. But beef cows can graze these lands as well and this makes up another area for comparison which may be more important in other Basin and Range states than in Utah and Nevada. In addition, there are private pasture lands that are used specifically for grazing. We will take up this in connection with Appendix I.

#### Forest Service Reports on AUMs for Selected States (Appendix H)

The Forest Service AUMs are reported in Annual Reports similar to the BLM reports. For Appendix H we accessed these reports. As for the BLM numbers by state, these numbers are reported in a list and then graphed for ease of comparison to the BLM numbers. In one of the charts we have plotted the "Permitted" and "Paid" AUMs and in a second chart we have plotted beef cattle numbers from NASS against the AUMs for each state. Again, there is room for confusion. In some states there is a rather close coherence between cattle numbers and AUMs divided by months, while in others there is not a complete coherence in the two numbers. The reason may be due to private grazing in some of the included states.

The obvious change in counting of AUMs is to reduce them to comparisons to beef cattle numbers. These comparisons would be easier if, like the Forest Service, the BLM consistently reported numbers of animals grazed per state. The combination of the two agencies' cattle numbers grazed could then be compared to NASS beef cow numbers and a clearer picture of the grazing patterns would emerge. Obviously we need additional data on the states to fully make the AUM and beef cattle numbers more consistent. This would give ranchers and others interested in agriculture a sounder basis for participating in discussions relating to grazing allocations.

## **7. Results of Input Output Analysis**

The Input output analysis was carried out only a selected county in Nevada and the State of Nevada. This was because the input output models were not available for other states in the Region. We

calculated economic benefits in the input output models to find the value of cattle grazing for two reasons; the calculation of economic impacts of cattle grazing at State and county level. For the State, the economic benefits are much smaller than for the agricultural county, Elko. The calculation of implications of increased cattle grazing used 1982 Census estimates for Elko County. Of course, this estimate has major implications for the wildfire model and related concerns about the economics of increased cattle numbers, causes and costs of wildfire damage.

A review of the IMPLAN model

To calculate economic impacts, an input-output framework was used. Input-output analysis uses a transaction matrix to show how outputs from one industry become inputs for another. The columns of the matrix represent purchases made by each industry and the rows represent sales made by each industry. This framework facilitates tracing of supply chains between industries by calculating proportions of industry sales and purchases to every industry represented in the input-output matrix.

For example, consider a specific element ( $x_{ij}$ ) of the transaction matrix, where  $i$  is the output (selling) industry and  $j$  is the input (purchasing) industry. This element in the matrix indicates the quantity of output of industry  $i$  sold to industry  $j$ ; or conversely, the quantity of input purchased by industry  $j$  from industry  $i$ . From this "base" transaction matrix, a second matrix of direct requirements can be derived by dividing each element of the transaction matrix by the sum of its column:

$$a_{ij} = \frac{x_{ij}}{\sum_{i=1}^i x_{ij}}$$

This yields a matrix of backward linkages, direct requirements that indicate the proportion of inputs from each industry  $i$  needed to produce one unit of output by industry  $j$ . Subtracting this matrix from an identity matrix and inverting the resulting matrix  $[(I - A)^{-1}]$  yields a matrix of final demand coefficients. The sum of the columns of the final demand matrix yields the output multiplier.

The analysis was conducted using the software, IMPLAN, which automates the input-output method. IMPLAN defines industries using a "proprietary classification system" that separates industries into 536 categories and assigns each an industry code numbered, 1 through 536. In general, IMPLAN industry codes correspond to those defined by the North American Industry Classification System (NAICS) at varying levels of specificity. Beef cattle ranches fit under the IMPLAN industry code 11, "Beef cattle ranching and farming, including feedlots and dual-purpose ranching and farming", which was used to calculate the economic impact for the region.

An "analysis-by-parts" method is used to calculate impacts for Elko County individually, and Nevada as a whole. With this method, ranch level budget data is used to derive percentages of expenditures of beef cattle ranches on commodities produced by other industries and on employee compensation and proprietor income. Impacts are determined by inputting the spending percentages with an accompanying spending level, and the level of employee compensation and proprietor income. IMPLAN uses these values to calculate Direct, Indirect, Induced, and Total economic impacts.

Direct effects are the initial changes in demand in the economy caused by the activity of green jobs. Indirect effects are changes in the same and other industries caused by the changes in inter-industry

purchases through the supply chain of the green sectors. Induced effects are changes in consumption caused by changes in employee compensation from both direct and indirect effects. The total contribution effect is the sum of direct, indirect, and induced effects.

## Results

Nevada has seen a decline in the number of cattle over the last few decades. In 1982, there were 359,000 beef cattle in the state, whereas today and averaging over the years of 2014 to 2016; the number of cattle in Nevada has decreased 223,035. Elko County has by far the greatest portion of cattle ranches in the state and has seen a decline from 101,578 head in 1982 to 72,214 head that is an average of 2014 to 2016 numbers.

Impacts for Elko County are presented in Tables 2 and 3. Table 2 shows the current impacts of beef cattle in Nevada, given the current number of cattle in the County. Table 3 shows the hypothetical impacts of beef cattle ranch operations, if the number of cattle in the County were returned to the numbers of 1982. The difference in impacts in Tables 2 and 3 represent an increase of 29,364 cattle in the county.

**Table 2: Elko County 2016 with 2014-2016 Average Number of Cattle**

Impact Type	Employment	Labor Income	Total Value Added	Output
Direct Effect	188.9	5,990,868	8,802,269	67,709,764
Indirect Effect	259.6	14,673,361	11,063,141	20,883,959
Induced Effect	55.6	2,101,932	3,789,814	6,187,388
Total Effect	504.1	22,766,161	23,655,225	94,781,110

**Table 3. Elko County with 1982 Number of Cattle**

Impact Type	Employment	Labor Income	Total Value Added	Output
Direct Effect	265.7	8,426,904	12,381,490	95,242,230
Indirect Effect	365.1	20,639,910	15,561,687	29,375,890
Induced Effect	78.2	2,956,630	5,330,846	8,703,332
Total Effect	709.1	32,023,444	33,274,023	133,321,452

According to the analysis, the current lower number of cattle is associated with approximately 200 fewer jobs in Elko County; \$10 million less labor income, \$10 million less GDP, and \$40 million lower value of output.

Impacts for the entire state of Nevada are presented in Tables 4 and 5 and are analogous to Tables 2 and 3. The direct and related effects are reported in Table 4. The differences in the impacts are presented in Tables 4 and 5 represent an increase of 135,965 cattle in the state.

**Table 4: Nevada 2016 with 2014-2016 Average Number of Cattle**

Impact Type	Employment	Labor Income	Total Value Added	Output
Direct Effect	683.4	20,919,990	28,143,504	216,488,489
Indirect Effect	2,140.7	49,867,855	113,240,047	198,604,918
Induced Effect	305.7	12,055,132	23,464,118	37,509,411
Total Effect	3,129.9	82,842,977	164,847,668	452,602,818

**Table 5: Nevada 2016 with 1982 Number of Cattle**

Impact Type	Employment	Labor Income	Total Value Added	Output
Direct Effect	1,100.1	33,673,085	45,300,145	348,462,652
Indirect Effect	3,445.8	80,267,937	182,272,628	319,677,015
Induced Effect	492.1	19,404,095	37,768,144	60,375,628
Total Effect	5,038.0	133,345,117	265,340,917	728,515,295

The analysis shows that the current lower number of cattle is associated with approximately 1,900 fewer jobs in Nevada, \$51 million less labor income, \$100 million less GDP, and numbers for comparison, \$276 less value of output.

Taken together, the rest of the counties in addition to Elko would have approximately 1,700 fewer jobs, \$41 million less labor income, \$90 million less GDP, and \$236 million lower value of output due to the smaller number of cattle between 1982 and the average of 2014 – 2016.

These results show two things: First, the difference between county and state level evaluations. Rural counties have much greater impact of cattle grazing. Secondly, the results show bias that is implied by taking only state level information for evaluating the impacts of grazing cattle.

## **8. The Benefits of Reduced Wildfire Return Intervals from Novel Grazing Management**

This section examines the potential for the novel grazing management techniques (henceforth *novel grazing management*) considered in this report to reduce wildfire activity and suppression costs in two ecological states in the Wyoming Sagebrush Steppe ecosystem in the Great Basin. The analysis assumes that novel grazing management will reduce the size and frequency of rangeland wildfires, thereby reducing the wildfire return interval on the landscape. Given that we do not yet have empirical estimates of the impacts of novel grazing management on wildfire return intervals, this section reports from the results from a sensitivity analysis that explores what the impact of these novel grazing techniques would be if they resulted in a 10%, 25%, etc. increase in wildfire return intervals.

The analysis in this section uses the simulation presented in Taylor et al. (2013). The description of the model and its parameterization presented here reproduces information reported in Taylor et al. For information on the technical details of the simulation procedure, please refer to Taylor et al.

## Materials and Methods

This analysis considers the potential benefits of novel grazing management in terms of reduce wildfire activity and suppression costs in the Wyoming Sagebrush Steppe (WSS) ecosystem in the Great Basin. The WSS ecosystem is generally found between 4,700 and 6,500 feet and comprises roughly 37.8 million acres in the Great Basin (26% of the 145-million-acre Great Basin). Figure 3 depicts the geographic extent of the WSS ecosystem in the Great Basin. The spread of exotic annual grasses such as cheatgrass (*Bromus tectorum*) is most prominent resource management concern relative to wildfire in the WSS ecosystem (Pellant, 1994; Miller and Tausch, 2001).

The simulation model captures rangeland ecosystem dynamics using the state-and-transition model (STM) framework from rangeland ecology (Stringham et al., 2003). The stylized STM of the WSS ecosystem used in the simulations is illustrated in Figure 4. The STM consists of three ecological states: 1. Perennial grasses and sagebrush with a small presence of invasive annual grasses characterize the “healthiest” state (WSS-1); 2. Overgrown “decadent” sagebrush with reduced perennial grasses and increased annual grasses (WSS-2); and 3. Invasive grass dominated state (WSS-3). Novel grazing management is assumed to be applied in WSS-2 and WSS-3, i.e., in the two ecological states where invasive annual grasses are present.

The wildfire suppression costs data comes from a data set of 400 wildfires occurring from 1995 through 2007 in USFS Region 4, the Intermountain Region (which includes Wyoming, Utah, Idaho, Nevada, and portions of Colorado and California). Wildfire suppression costs are reported in Table 6. See Taylor et al. (2013) for details.

**Table 6. Wildfire Suppression Costs (\$000 in 2010 dollars; 000’s of acres)**

Ecological State	NFDRS Fuel Model <sup>a</sup>	No. Obs	Avg. \$/fire	Total expenditure	Avg. Acres/fire	Total Acres burned	Avg. \$/acre
WSS-1	T and L	43	\$441.9	\$19,003.7	2.3	100.0	\$190.1
WSS-2	B	14	\$844.1	\$11,817.6	1.1	15.0	\$788.7
WSS-3	A	12	\$1,314.8	\$15,777.5	12.9	154.2	\$102.3


<sup>a</sup> See Taylor et al. (2013) for discussion of National Fire Danger Rating System (NFDRS) fuel models.

Wildfire is modeled as a stochastic event that may or may not occur in a given year. Wildfire return intervals for each state in the WSS ecosystem are taken from the “Wyoming Sagebrush Steppe” LANDFIRE Rapid Assessment Vegetation Model. Wildfire return intervals are reported in Table 7. See Taylor et al. (2013) for details.

**Table 7. Wildfire Frequency**


	Ecological State

	WSS-1	WSS-2	WSS-3
Wildfire-return interval (years)	107	75	9
Annual large fire probability	0.009	0.013	0.111



### Legend

-  Great Basin Boundary
-  Wyoming Sagebrush
-  Mountain Big Sagebrush




**Figure 3. Geographic Distribution of Wyoming Sagebrush Steppe (WSS) Plant Communities in the Great Basin. (Reproduced from Taylor et al. (2013).)**

The number of years to transition from an ecological state to another through ecological succession absent disturbance (i.e., wildfire) were taken from the “Wyoming Sagebrush Steppe” LANDFIRE Rapid Assessment Vegetation Model. Transition between ecological states after wildfire depend on the presence of annual grasses in the understory. Wildfire in WSS-1 is ecologically beneficial and restores the system to an earlier successional stage in WSS-1. Wildfire in WSS-2, however, cause the system to transition to WSS-3. See Taylor et al. (2013) for details. Transitions with and without wildfire for the WSS ecosystem are summarized in Table 8.

**Table 8. Transitions between States**

	Ecological State		
	WSS-1	WSS-2	WSS-3
Time to Transition w/o Wildfire	60 years → WSS-2	NA	NA
Transition with Fire	→ Year 1 in WSS-1	→ WSS-3	Stay in WSS-3



**Figure 4. Wyoming Sagebrush Steppe Stylized State-and-Transition Model (Reproduced from Taylor et al. (2013).)**

**Results**

Tables 9, 10, and 11. the reported results on the potential benefits of the novel grazing management if it is successful at increasing the wildfire return interval on the landscape, thereby making wildfires less frequent. Tables 4, 5, and 6 report results for five hypothetical increases in the wildfire return interval: 25%, 50%, 100%, 150%, and 200%. The results assume that the novel grazing treatment is applied in both WSS-2 and WSS-3.

Table 9 reports results when the initial state is WSS-2 and WSS-3. The results indicate that if novel grazing management is successful at increasing the wildfire return internals, it can result is substantial reductions in public wildfire suppression costs. For example, if novel grazing management were to double the wildfire return interval, it would lead to 1.13 fewer fires over a 50-year time horizon, on

average, when the initial state is WSS-2. The expected net present value of the suppression cost saving from this reduction in wildfire activity is \$112 per acre. When the initial state is WSS-3, novel grazing management is predicted to lead to 2.69 fewer wildfires and \$142 in per acre suppression costs savings over 50 years. Per acre suppression costs do not increase linearly with wildfire occurrence when the initial state is changed from WSS-2 to WSS-3 because wildfires in WSS-3 (i.e., on annual grass dominated rangelands) are more likely to be allowed to burn without active suppression efforts, so that the pre-acre suppression expenditures are lower in WSS-3 compared to WSS-1 or WSS-2.

**Table 9. Grazing Treatment (\$ per acre; 2010 dollars)**

	Fire Return Interval Increase from Treatment					
<b>Initial State = WSS-2</b>	<b>Baseline</b>	<b>25%</b>	<b>50%</b>	<b>100%</b>	<b>150%</b>	<b>200%</b>
Mean Number of Wildfires <sup>a</sup>	1.92	1.40	1.09	0.79	0.48	0.34
Reduction in Number of Wildfires	-	0.52	0.83	1.13	1.44	1.58
Mean Total Suppression Costs <sup>b</sup> (NPV)	\$277.0	\$225.4	\$221.2	\$165.3	\$101.5	\$77.0
Total Cost Savings	-	\$51.6	\$55.8	\$111.7	\$175.5	\$200.0
Mean Annual Suppression Costs	\$10.4	\$8.6	\$7.7	\$6.1	\$4.0	\$3.0
Annual Cost Savings	-	\$1.9	\$2.8	\$4.3	\$6.4	\$7.4
Final State <sup>c</sup> (WSS-1, WSS-2, WSS-3)	(0,527,473)	(0,594,406)	(0,634,366)	(0,688,312)	(0,777,223)	(0,822,178)
<b>Initial State = WSS-3</b>	<b>Baseline</b>	<b>25%</b>	<b>50%</b>	<b>100%</b>	<b>150%</b>	<b>200%</b>
Mean Number of Wildfires	5.54	4.44	3.76	2.85	2.16	1.87
Reduction in Number of Wildfires	-	1.10	1.78	2.69	3.38	3.67
Mean Total Suppression Costs (NPV)	\$299.2	\$242.9	\$207.1	\$156.7	\$115.4	\$104.9
Total Cost Savings	-	\$56.3	\$92.1	\$142.4	\$183.8	\$194.2
Mean Annual Suppression Costs	\$11.2	\$9.1	\$7.7	\$5.8	\$4.4	\$4.0
Annual Cost Savings	-	\$2.1	\$3.4	\$5.3	\$6.8	\$7.2

<sup>a</sup> The simulation model considers the progression of the WSS ecosystem over 50 years

<sup>b</sup> All dollar values are discounted at a constant rate of 3%.

<sup>c</sup> Final State' is the final state of the system (WSS-1, WSS-2, or WSS-3) after 50 years. Final state results not reported when the initial state is WSS-3 (the annual grass dominated state) because the system never exits WSS-3 in this state.

An additional benefit of novel grazing management in WSS-2 is that by reducing wildfire, the system is less likely to transition to WSS-3 as a result of wildfire. For example, when the initial state is WSS-2, the system ends up in WSS-3 after 50 years in 47% of simulation runs under our baseline assumptions. In contrast, if the grazing treatment successfully double the fire return interval, the system ends up in WSS-3 in only 37% of simulation runs.

Novel grazing management increases the costs of herd management. In economic terms, these are variable (as opposed to fixed) costs. Tables 9, 10, and 11 report results for the predicted annual per-acre suppression cost savings. These cost savings, scaled up to the size of the ranch, can be compared to the increased cost herd management over a grazing season and, as such, can be used to evaluate whether a public program subsidizing ranchers to pursue this novel grazing strategy can be justified on the basis of wildfire suppression cost savings alone. Such a program could potentially be implemented through a Farm Bill program such as Natural Resources Conservation Service’s *Environmental Quality Incentives Program*. The results in Table 4 indicate that the expected annual per-acre suppression costs savings range from \$1.9-\$7.4 when the initial state is WWS-2 and \$2.1 to \$7.2 when the initial state is WSS-3 depending on the anticipated increase in wildfire return intervals in WSS-2 and WSS-3.

Table 10 reports the results when in addition to reducing wildfire frequency, novel grazing management alters plant community composition in a way that enhances ecosystem resilience and encourages post-fire recovery. Specifically, the results reported in Table 10 assumes that wildfires in WSS-2 are ecological beneficial and cause the system to transition to WSS-1. In contrast, the results reported in Table 4 assume that wildfires in WSS-2 cause the system to transition to WSS-3. Table 5 shows that if novel grazing management enhances ecological resilience it will lead to a further reduction in wildfire activity and suppression costs. For example, comparing the Table 4 and 5 results for a 100% increase in wildfire return interval, including ecological resilience leads to 1.55 fewer fires over 50 years versus 1.13 with a \$141 in per acre suppression cost savings versus \$112. These results indicate that the economic benefits of novel grazing management will depend critically on whether it increases the ecological resilience treated plant communities.

**Table 10. Grazing Treatment: Ecological Resilience (\$ per acre; 2010 dollars)**

Initial State = WSS-2	Fire Return Interval Increase from Treatment					
	Baseline	25%	50%	100%	150%	200%
Mean Number of Wildfires <sup>a</sup>	1.92	0.54	0.45	0.37	0.27	0.22
Reduction in Number of Wildfires	-	1.38	1.48	1.55	1.65	1.70
Mean Total Suppression Costs <sup>b</sup> (NPV)	\$277.0	\$207.5	\$171.1	\$136.1	\$98.4	\$77.1
Total Cost Savings	-	\$69.5	\$105.9	\$140.9	\$178.6	\$199.9
Mean Annual Suppression Costs	\$10.4	\$7.6	\$6.4	\$5.0	\$3.7	\$2.9
Annual Cost Savings	-	\$2.8	\$4.0	\$5.5	\$6.7	\$7.5

Final State <sup>c</sup> (WSS-1, WSS-2, WSS-3)	(444,556,0)	(424,576,0)	(358,642,0)	(291,709,0)	(216,784,0)	(178,822,0)
---	-------------	-------------	-------------	-------------	-------------	-------------

<sup>a</sup> The simulation model considers the progression of the WSS ecosystem over 50 years

<sup>b</sup> All dollar values are discounted at a constant rate of 3%.

<sup>c</sup> Final State<sup>c</sup> is the final state of the system (WSS-1, WSS-2, or WSS-3) after 50 years.

Table 10 also indicates that the primary benefit of enhanced ecological resilience may not be from reductions in wildfire activity or suppression costs, but rather from preventing the system from moving to WSS-3, which entails permanent degraded resource values. In our simulations, when the initial state is WSS-2, allowing novel grazing management to enhance ecological resilience causes the system to be restored to WSS-1 in the majority of simulations, and prevents the system from transitioning to WSS-3. Note that Table 5 only considers when the initial state is WSS-2. The analysis assumes that, on its own, novel grazing management is not sufficient to rehabilitate the system from WSS-3.

Table 11 reports results from an alternative baseline where future fire return intervals are lower than historic averages as a result of invasive plants, changes in disturbance regimes, climate change, and other factors. These reductions in wildfire return intervals are anticipated in the WSS ecosystem and in other ecosystems throughout the Great Basin (Baker, 2009; Romme et al., 2009). Table 6 indicates that the benefits of novel grazing management are likely to increase significantly if wildfires are more frequent in the future. This result suggests that evaluating novel grazing management using current estimates of wildfire return intervals may significantly understate its potential benefits.

**Table 11. Grazing Treatment: Shortened Fire Return Intervals (\$ per acre; 2010 dollars)**

Initial State = WSS-2	Fire Return Interval Increase from Treatment					
	Baseline	25%	50%	100%	150%	200%
Mean Number of Wildfires <sup>a</sup>	5.74	3.99	3.10	2.03	1.48	1.07
Reduction in Number of Wildfires	-	1.75	2.64	3.71	4.26	4.66
Mean Total Suppression Costs <sup>b</sup> (NPV)	\$568.8	\$464.4	\$390.1	\$281.8	\$236.3	\$185.1
Total Cost Savings	-	\$104.4	\$178.7	\$287.0	\$332.5	\$383.8
Mean Annual Suppression Costs	\$21.6	\$17.5	\$15.1	\$10.4	\$8.6	\$7.1
Annual Cost Savings	-	\$4.1	\$6.5	\$11.2	\$12.9	\$14.5
Final State <sup>c</sup> (WSS-1, WSS-2, WSS-3)	(0,255,745)	(0,353,647)	(0,398,602)	(0,534,466)	(0,580,420)	(0,647,353)
Initial State = WSS-3	Baseline	25%	50%	100%	150%	200%
Mean Number of Wildfires	11.0	8.8	7.3	5.5	4.5	3.7

Reduction in Number of Wildfires	-	2.2	3.7	5.5	6.5	7.3
Mean Total Suppression Costs (NPV)	\$588.8	\$475.0	\$386.6	\$303.1	\$248.3	\$201.7
Total Cost Savings	-	\$113.8	\$202.2	\$285.8	\$340.5	\$387.1
Mean Annual Suppression Costs	\$22.3	\$17.9	\$14.8	\$11.4	\$9.3	\$7.6
Annual Cost Savings	-	\$4.4	\$7.5	\$10.8	\$13.0	\$14.7

<sup>a</sup> The simulation model considers the progression of the WSS ecosystem over 50 years

<sup>b</sup> All dollar values are discounted at a constant rate of 3%.

<sup>c</sup> Final State' is the final state of the system (WSS-1, WSS-2, or WSS-3) after 50 years. Final state results not reported when the initial state is WSS-3 (the annual grass dominated state) because the system never exits WWS-3 in this state.

## Conclusions

- If novel grazing management is successful at increasing the wildfire return internals (i.e., reducing wildfire frequency), it can result in substantial reductions in wildfire suppression costs, with the net present value of suppression cost savings over a 50-year horizon ranging from a low of around \$50 per acre to over \$300 per acre depending on model assumptions.
- The economic benefits of novel grazing management will depend critically on whether it increases the ecological resilience of treated plant communities.
- Novel grazing management will make WSS ecosystems less likely to transition to an annual grass dominated state, thereby avoiding permanently degraded resource values.
- The benefits of novel grazing management are significantly greater if future wildfire return intervals are reduced due to climate change and other factors identified in the literature. This suggests that evaluating novel grazing management using current estimates of wildfire return intervals may significantly underestimate its potential benefit.

## Limitations

- The results reported here provide conservative estimates of the benefits of novel grazing management. Most significantly, the analysis only considers wildfire suppression cost savings. Other potential benefits not considered in the analysis include improved forage for livestock and economic outcomes for ranchers, reductions in wildfire damages to private property and public infrastructure, and improvements in wildlife habitat, recreation opportunities, erosion control, and other ecosystem goods and services.
- The analysis does not consider the cost of novel grazing management. As such, the analysis can only answer the more limited question of at what per acre costs will novel grazing management be justified on the basis of wildfire suppression cost savings alone. The results reported here indicate that a public subsidy for ranchers implementing novel grazing management

exceeding 10\$ per acre per year could be justified. This would imply a negative grazing fee for ranchers implementing novel grazing management on public rangelands.

#### Impact of Extended Grazing of Cheatgrass on a Cow-calf Budget: An Example

Using grazing prescriptions that include an estimated two months of fall cheatgrass not only has environmental benefits, but also directly impacts the costs and profitability of cattle ranches. Substituting two months of grazing for two months of much costlier winter feeding of hay and grain can add a significant amount to ranchers' bottom lines. We used the Elko County 700 cow-calf operation 2014-2016 average budget to estimate such impacts. A number of simplifying assumptions were made for this exercise, and the results should be viewed as a general example. Assumptions include:


1. The original grazing season is six months and would be expanded to eight months.
2. The original winter feeding period is six months and would be reduced to four months.
3. There is adequate cheatgrass area for all cattle on the ranch during the expanded grazing season.
4. We did not adjust other variable costs such as changing protein and other supplements, transportation costs or labor costs. We realize that these might be different but would require meeting with a rancher panel to discuss.
5. Adequate nutrition was obtained during the extended grazing system and animal weights are not affected.

Figure 5 illustrates the estimated difference between the traditional grazing schedule and one that included an extended period of cheatgrass grazing. Grazing fees per cow would increase approximately \$4, more than twice the \$1.35 per AUM, as again, our assumption is that all cattle utilize extended grazing, and are included in per cow averages.

#### Cheatgrass Fuel Load Reduction and Water Availability

In addition, the extended grazing on cheatgrass would require protein supplements for proper nutrition. Assuming one pound per head per day, a total of 60 pounds per head would be required for the two months additional grazing. Again, we assume that all cattle would require the supplement, so we approximate the total used to be somewhere in the neighborhood of 60,000 pounds. Recent prices for 24% protein beef cattle supplements for pasture and forage fed cattle run around \$0.33 per pound, or a total expense of around \$19,800. Allocating that to a 700 cow herd gives a price per cow of around \$28. Although grazing fees show a modest increase, and protein supplement costs are substantial, hay and

grain costs, the largest cost category, decline by one-third, or \$91 per cow. With the added protein supplement costs, and the small increase in grazing costs, the overall variable costs will be reduced by \$61 per cow. With cattle prices being the same with both treatments, this translates directly into increased profits of the same magnitude. Again, it must be emphasized that this is a simplified view and shows the maximum of savings and increased net returns.



**Figure 5. Comparison of Costs and Net Returns from Traditional and Extended Grazing**

For the 700 cow ranch, this translates into approximately \$42,000 higher net returns for the 2014-2016 average period. If adequate cheatgrass existed in the appropriate locations that all ranchers in Elko County could utilize the extended grazing period, the county-wide impacts could be as much as \$4.2 million. However, we must not assume that this would be a persistent possibility. Reducing the cheatgrass load in areas that could subsequently move to improved vegetation stages would eventually eliminate much of the cheatgrass. On the other hand, there could also be an increase in native species and expanded grazing areas that could accommodate higher numbers of cattle during traditional grazing seasons. Such decisions on grazing allotments and permits would ultimately be the purview of the BLM and Forest Service.

In addition, the removal of cheatgrass has implications for fuel reduction. Cattle grazing cheatgrass will consume from 20 to 25 pounds per day for the two fall months. We select 25 pounds as the figure because the cattle are rather mature during the fall season. At the upper limit of 25 pounds times 60 days this cheatgrass consumption level equals 1,500 pounds per head for the fall season. Multiplying

this number times the approximate number of cattle available for fall grazing which is approximately 70,000 in Elko County equals an estimated 1.05 million pounds of cheatgrass consumed during the fall two-month season. A natural comparison would be to relate this number to the annual growth of cheatgrass in Elko County, which would give a number that would help the BLM evaluate the impact of cattle fall grazing.

Cheatgrass" thatch" left from earlier years is a good medium for the germination of cheatgrass. When the thatch is consumed by cattle in the fall, the rate of germination will decrease and there will be fewer plants that germinate the following year due to greater exposure to weather conditions. Depending on conclusive evidence of germination of cheatgrass in thatch, this factor may be a major contribution to cheatgrass control due to lower germination in the spring when the growing season for cheatgrass commences.

There are of course limitations to this fall grazing strategy—water must be available for cattle which is problematic in some cheatgrass infested areas. There must be adequate water in the proximity of the protein supplement to make both readily accessible to the herd. As well, there is the commitment of ranch hands to the change locations of cattle for full implementations of fall grazing. One of the reasons for the avocation of fall grazing is that the cattle do not stray far from the protein feeder that dispenses the nutrients that supplement the carbohydrates in the cheatgrass. Cattle will not go much more than 300 yards from the protein feeder, making it possible to graze the cattle with limited ranch hand supervision. All that is needed is to pull the feeder along as the cheatgrass is consumed.

Finally, there is some amount of grazing of perennials during the fall grazing season. If the grazing of cheatgrass is in areas that have near 100 percent infestation, there is not much grazing of perennials. However, if the cheatgrass is mixed with perennials in cheatgrass infested areas there will be some grazing of perennials. Some estimates of the grazing of perennials put the level at about 20 percent. This is not problematic for the perennials since they are in a dormant state during the fall season after a hard frost.

## **9.**

### **Alternative Approaches to Scoring Rules**

#### Review of the Scoring Literature

The "scoring" literature is extensive, often provided under the title "environmental scoring". This reflects a theme from the beginning of the approach which featured the environmental dimension. Scoring can now be applied to firms or organizations with profit incentives, communities (the approach we will

select) and regions or nations. The best example of scoring applied to communities is from Wikipedia (Wikipedia, 2018). It lays out the approach in simple terms and it is segmented into four areas; environmental concerns, social concerns, corporate governance concerns and responsible investment. These will be demonstrated to provide a scope that is sufficient to cover the communities in the Basin and Range Region. Again, the motivation for the communities approach is to eventually test the mechanism in several communities in the Basin and Range Region.

## History

The history of community involvement in scoring and other involvements in the future of the establishments is rather recent. Initially, Milton Friedman was an advocate of not tying the hands of communities to eliminate “soft” concerns about the future of the communities focusing only on profitability. This was a central theme of early approaches to scoring. The approach emphasized that concerns about the community were drags on the profitability of firms and should not be considered if the objective was to grow and develop.

This began to lose ground as other leaders were more interested in the community at large than in single firms. A major departure from the older philosophy came with a writing of Coleman on “Social Capital in the Creation of Human Capital” (Coleman, 1988). His assertion was that for the long run good of society other factors should be included in scoring than simply profitability. This article led to a proliferation of other writings on the subject and to a new concept of community leaders as participants in the management of community activities.

The evolution of scoring was a later extension of more involvement in national and international scoring and in the future of communities. Currently there are many focuses on aspects of communities ranging from those with an environmental bent to those with particular emphasis on problem areas for communities. In this summary we are contemplating using the scoring approach to deal with agricultural communities that are threatened by wildfire and dealing with fuel loads that are a byproduct of cheatgrass domination in the local area.

## Procedure

The procedure is discussed in several documents and relates to first going to the communities and discussing the approach with citizens’ groups reflecting a cross section of interests in the communities- usually 15 to 20 members. The idea is to go through the scoring system and solicit community members’ input on the issues that they will be asked to score. There will likely be broad discussion of the questions or assertions that they are asked to score, and the system that will be applied for summarizing the scores. This process should take about 2 – 3 hours.

After several weeks the organizers of the scoring exercise should return to the communities for an application of the scoring method. This will likely be preceded by a number of questions based on the earlier visit. After answering about process and responding to these questions, the community leaders will proceed to answer the questions on issues designated in the scoring sheet. The scoring sheet must be rather brief since the community participants will be interested in returning to their occupations.

With this preliminary review of the process we will turn to the concerns raised in the Wikipedia review of the approach. Here we will suggest some of the questions or issues that the community participants may be asked to respond to.

For environmental concerns the list would likely include:

#### Climate change

The question or issue for this subject will be whether or not the members of the community believe that there has or will have the prospects that it will impact the part of the community that is important to them

#### Sustainability

The issue is if the community is depleting natural resources that are critical to the success of the community for development or growth. The issue may be more appropriately discussed in terms of efficiency or utilizing available resources.

#### Social concerns

##### Diversity

The level of inclusion of all citizens in recruitment policies broadening the pool of talent.

##### Consumer protection

The right of consumers to be considered in “buyer beware” issues related to purchase of commodities or purchases of land or other fixed assets.

## Animal Welfare

Concerns about animals for inputs to the food system or for companions.

## Corporate Governance

### Management Structure

Are employees managed in responsible ways by management?

### Employee relations

Are employees effectively handled.

### Executive compensation

Are executives compensated for their efforts?

## Investment Strategies

### Positive selection

Selection of investors that is regarded as inclusive.

### Activism strategies

Seeking out investors that are not forthcoming

### Engagement

Engagement of all of the community.

Consultation roles

Using consultants effectively.

Exclusion

Exclusion in any aspect of investment is a no/no.

Integration

Effectively integrating all of the community.

All of these concerns will be pointed toward grazing issues in the agricultural communities. And the grazing of cheatgrass will be a major concern along with the degradation of range lands which appears to be a major concern of the BLM. Degradation can be reversed but should be a part of any community scoring exercise. The focus in fact will be to energize the communities to deal with fuel loads and their build up on the range land. The interests will be related to wildfires and their threat to community infrastructure. Many of these rural communities have little investment but are dependent for their basic income on cattle that are grown on the range land.

### Example Applications

There are a number of applications that may serve the practical applications of scoring. We review each of these briefly to suggest themes that should be perused in developing the applications of scoring to communities in the Basin and Range Region.

STAR communities is an approach that may be useful in addressing the community needs. It is a national system that ranks communities or sets of communities in terms of achieving their targets. It is a national system and applies to various community problems or issues (STAR Communities 2018).

Another treatise is on the tool box for applications of scoring to communities (Community Tool Box 2018). This tool box uses a logic model to track community responses and provide structure to the summary of scoring. The logic model has been applied in many systems like agricultural extension, contract management and other fields for providing a simple structure for managing the process and the scoring exercise.

The Eccles et al. paper provides a good survey of scoring exercises and how the results are internalized in communities. This is more of a through-going paper and makes many points about scoring. It is a paper that would be good for beginners in the field of scoring.

The Hokunui (2017) application is close to the approach that would be taken in dealing with agricultural rural communities in the Basin and Range Region. It has a focus on agriculture and deals with a multitude of issues related to the growth and development of agriculture. Some of the features in this study are related to the kinds of issues that would be front and center in the Basin and Range Region.

The Rose Marcario (2014) paper is from Chile and also related to agriculture and its place in rural communities. This is a comparatively long study but interesting to read for background on the scoring approach.

The Morhardt (2002) study uses ISO 14031 as a basis for making a study of an area or community. This, like the STAR communities is another template for addressing communities.

## **10.**

### **Conclusions and Implications**

We have investigated the costs and revenues for cow-calf operations within the Basin and Range region. Specifically, we updated budgets for operations of various sizes in each of the states within the region for the time period of 2014-2016 so that costs and revenues can be compared across counties and regions.

Using the budgets, we integrated county beef cow inventories to calculate gross costs and revenues, as well as net returns for specific counties and regions to arrive at estimates of the value of the cow-calf industry in the counties and regions. Table 1 in this report summarizes these county estimates for budget areas, states, and the Basin and Range region.

The estimates of revenues, costs, and returns, as well as cattle inventories were input into an input-output framework to estimate the contribution of the cow-calf industry on the Nevada and Elko County economies. These impacts are summarized in Tables 2 through 5. They illustrate the difference between the state level and county level relative impacts.

We further considered the effects of grazing cheatgrass within the region utilizing a fire model to reduce highly-flammable fuel loads and estimate savings on fire suppression costs that would occur with reduced wildfire occurrence. Grazing prescriptions that include fall cheatgrass utilization not only reduce the fuel loads on rangelands, it also offsets some winter feeding costs, especially for ranchers in more northern locations within the region.

We applied the cheatgrass grazing possibilities to Elko County to arrive at the impact of extended fall cheatgrass grazing on the revenues, costs, and returns for the county. The prospective quantity of cheatgrass utilized during the fall grazing season would amount to 25 pounds per head per day, or a total of 1,500 pounds for a two-month period. If all cattle in Elko County were able to be grazed on cheatgrass in the fall, we estimate that the reduction in cheatgrass would amount to 1.05 million pounds.

Lastly, we performed a preliminary investigation of scoring of cattle production, environmental conditions and decreased threats to communities by reducing fuel loads. These scoring techniques could be applied to regions smaller than the Basin and Range Region to illustrate benefits on a localized basis. The idea is to address alternatives for communities impacted by fire and alternative grazing prescriptions.

## 11. References

- Abatzoglou, J.T., C.A. Kolden. 2011. Climate change in western US deserts: potential for increased wildfire and invasive annual grasses. *Rangeland Ecology and Management* 64:471-478
- Baker, W.L. 2009. *Fire Ecology in Rocky Mountain Landscapes*. Washington, D.C.: Island Press.
- Balch, J.K., B.A. Bradley, C.M. D'antonio, J. Gómez-Dans. 2013. Introduced annual grass increases regional fire activity across the arid western USA (1980–2009). *Global Change Biology* 19:173-183.
- Bates, J.D., K.W. Davies. 2014. Cattle grazing and vegetation succession on burned sagebrush steppe. *Rangeland Ecology and Management* 67:412-422.
- Bonan, G.B. 1997. Effects of land use on the climate of the United States. *Climatic Change* 37:449-486.
- Chambers, J.C., B.A. Bradley, C.S. Brown, C. D'Antonio, M.J. Germino, J.B. Grace, S.P. Hardegree, R.F. Miller, D.A. Pyke. 2014. Resilience to stress and disturbance, and resistance to *Bromus tectorum* L. invasion in cold desert shrublands of western North America. *Ecosystems* 17:360-375.
- Chambers, J.C., B.A. Bradley, C.S. Brown, C. D'Antonio, M.J. Germino, J.B. Grace, S.P. Hardegree, R.F. Miller, D.A. Pyke. 2014. Resilience to stress and disturbance, and resistance to *Bromus tectorum* L. invasion in cold desert shrublands of western North America. *Ecosystems* 17:360-375.
- Coleman, James S., "Social Capital in the Creation of Human Capital". *American Journal of Sociology*, 94, 1988.
- Community Tool Box; a Logic Model or Theory of Change, google 2018
- Coupal, R. and D. Holland. 1995. "On the Use of Farm Budgets in Interindustry Analysis: An Example from Washington State Wheat Study". Washington State University, Department of Agricultural Economics, A.E. 95-10. Pullman, Washington.
- Clemets, Charles. D. and Dan Harmon. 2018. *Rangeland Rehabilitation Laboratory Report-USDA Great Basin Rangeland Research Unit USDA ARS*. Mimeo.
- Eccles, R. O., Ioannou Ioannou and George Serafeim, *The Impact of Sustainability on Organizational Processes and performance*, Unpublished manuscript, 2011
- Friedman, Milton and J. Rose, *Free to Choose, a Personal Statement*, Harcourt, 1980
- Global Green Economy Index (GGEI) 2016*.
- Hokunui Maui, Our Vision is to have a Thriving and Relationship between Community and Agriculture (<https://www.hokunui.com/>)
- Johnson, S. R. 2017. "An Economic Analysis of Cattle Numbers and Feed Sources in the USDA Basin and Range Region" application for funding from the Public Lands Endowment Trust.
- Jones, A. 2000. Effects of cattle grazing on North American arid ecosystems: a quantitative review. *Western North American Naturalist* 60:155-164.
- Kline, J., 2004. *Issues in Evaluating the Costs and Benefits of Fuel Treatments to Reduce Wildfire in the Nation's Forests*. Research Note PNW-RN-542. USDA Forest Service, Portland, OR.
- Knapp, P.A., P.T. Soulé. 1996. Vegetation change and the role of atmospheric CO<sub>2</sub> enrichment on a relict site in central Oregon: 1960–1994. *Annals of the Association of American Geographers* 86:387-411

- Knutson, K.C., D.A. Pyke, T.A. Wirth, R.S. Arkle, D.S. Pilliod, M.L. Brooks, J.C. Chambers, J.B. Grace. 2014. Long-term effects of seeding after wildfire on vegetation in Great Basin shrubland ecosystems. *Journal of Applied Ecology* 51:1414-1424.
- Ludwig, J.A., B.P. Wilcox, D.D. Breshears, D.J. Tongway, A.C. Imeson. 2005. Vegetation patches and runoff-erosion as interacting ecohydrological processes in semiarid landscapes. *Ecology* 86:288-297.
- Leger, E.A., E.K. Espeland, K.R. Merrill, S.E. Meyer. 2009. Genetic variation and local adaptation at a cheatgrass (*Bromus tectorum*) invasion edge in western Nevada. *Molecular Ecology* 18:4366-4379.
- Marcario, Rose, Patagonia; Environmental and Social Initiatives, Chile publication, 2014.
- Meinke, C. W., Knick, S. T., Pyke, D. A., 2009, A Spatial Model to Preauthorize Sagebrush Landscape in the Intermountain West (USA) for Restoration, *Restoration Ecology*, 17(5) 652 – 659.
- Melgoza, G., R.S. Nowak, R.J. Tausch. 1990. Soil water exploitation after fire: competition between *Bromus tectorum* (cheatgrass) and two native species. *Oecologia* 83:7-13.
- Miller, R.F. and R.J. Tausch. 2001. The role of fire in pinyon and juniper woodlands: A descriptive analysis. In: K.E.M. Galley and T.P. Wilson (Ed.) Proceedings of the invasive species workshop: The role of fire in the control and spread of invasive species.
- Morhardt, Emil, Sarah Baird and Kelly Freeman, Scoring Corporate Environmental and MSustainability Reports Using GRI ISO 14031 and Other Criteria, Wiley online Library, 2002
- Mote, P.W., S. Li, D.P. Lettenmaier, M. Xiao, R. Engel. 2018. Dramatic declines in snowpack in the western US. *Climate and Atmospheric Science* 1:2.
- Nusser, S., F. Breidt, W. Fuller. 1998. Design and estimation for investigating the dynamics of natural resources. *Ecological Applications* 8:234-245.
- Pellant, M. 1994. History and applications of the Intermountain greenstripping program. In: Proceedings—symposium on ecology and management of annual rangelands; 1992 May 18-22; Boise, ID. Gen. Tech. Rep. INT-GTR-313. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 63-68.
- Pellant, M., B. Abbey, S. Karl. 2004. Restoring the Great Basin desert, USA: integrating science, management, and people. *Environmental Monitoring and Assessment* 99:169-179.
- Pierson, F.B., C.J. Williams, S.P. Hardegree, M.A. Weltz, J.J. Stone, P.E. Clark. 2011. Fire, plant invasions, and erosion events on western rangelands. *Rangeland Ecology and Management* 64:439-449.
- Romme, William H., Craig D. Allen, John D. Bailey, William L. Baker, Brandon T. Bestelmeyer, Peter M. Brown, Karen S. Eisenhart, M. Lisa Floyd, David W. Huffman, Brian F. Jacobs, Richard F. Miller, Esteban H. Muldavin, Thomas W. Swetnam, Robin J. Tausch, and Peter J. Weisberg. 2009. "Historical and Modern Disturbance Regimes, Stand Structures, and Landscape Dynamics in Pinyon Juniper Vegetation of the Western United States. *Rangeland Ecology & Management*. 62 (3), 203-22.
- Schmelzer, L., Perryman, B., Bruce, B., Schultz, B. McAdoo, K., McCain, G., Swanson, S., Wilke, j., Conley, K., 2014, Case Study: Reducing Cheatgrass (*Bromus Tectorum*) Fuel Loads Using Fall Cattle Grazing, *The Professional Animal Scientist* (30) 270 – 278.
- STAR Communities; new milestones in Urban Sustainability Movement, STAR Communities, google 2018
- Strachan, S., F. Biondi, J. Leising. 2011. 550-Year reconstruction of streamflow variability in Spring Valley, Nevada. *Journal of Water Resources Planning and Management* 138:326-333.

- Svejcar, T. and Perryman, B. L. 2014. Western Land Managers Will Need all Available Tools for Adapting to Climate Change, Including Grazing: A Critique of Beschta et al. *Environmental Management*. 30, 135 – 1038.
- Taylor, Michael H., Kimberly Rollins, Mimako Kobayashi, and Robin Tausch, 2013. The Economics of Fuel Management: Wildlife, Invasive Plants, and Dynamics of Sagebrush Rangelands in the Western United States. *Journal of Environmental Management* (28) 157 – 173.
- Trowbridge, W. Albright, T., Ferguson, S., Jun, J., Perryman, B., and Nowak, S., 2010, Explaining Patterns of Species Dominance in the Shrub Steppe Systems of the Junggar Basin (China) and the Great Basin (USA), *Journal of Arid Land*, 10 , 4033.
- University of Nevada, Reno, College of Agriculture, Biotechnology and Natural Resources. 2017. *Sustainable Dryland Agriculture*
- Vermeire, L.T., J.L. Crowder, D.B. Wester. 2014. Semiarid rangeland is resilient to summer fire and
- Wikipedia, Environmental, Social, and Corporate Governance, 2018
- Willis, D. and D. Holland. 1996. "Translating Farm Enterprise Budgets into Input-Output Accounts: Another Example from Washington State". Department of Agricultural Economics, Washington State University, Agricultural Economics Departmental Publication, Pullman, Washington.
- Yale Center for Environmental Law & Policy, and Center for International Earth Science Information Network at Columbia University. 2008, "*Environmental Performance Index*".
- Young, J.A., R.A. Evans, R.E. Eckert Jr, B.L. Kay. 1987. Cheatgrass. *Rangelands* 9:266-270.